


# ATLAS NSW Trigger



Alex Tuna

Harvard University

DPF @ Fermilab  
31 July 2017




# related talks this week

Yu: ATLAS MDTs for HL-LHC	Monday afternoon
Catrin: ATLAS trigger algs in Run 2	Monday afternoon
Heather: ATLAS trigger menu in Run 2	Monday afternoon
Tom: ATLAS Muon for HL-LHC	Wednesday afternoon
Xiangting: ATLAS tRPCs for Run 3+	Thursday morning
<b>Liang: ATLAS NSW Electronics</b>	<b>Thursday morning</b>
<b>Jared: CMS GEM for Run 3+</b>	<b>Thursday afternoon</b>

# introduction



Lake Geneva



LHCb

ATLAS


ALICE

CMS

LHC



# a 20-year plan for the LHC




instantaneous luminosity increasing — experiments must adapt!

ATLAS has suite of upgrades for near-term (Phase 1)  
and long-term (Phase 2)


# the new Small Wheel (NSW)

NSW replaces current SW —  
under construction now


comprised of eight layers each  
of two new detectors for ATLAS



*small Thin Gap Chambers (sTGC)*



*Micromegas (MM)*





# the new Small Wheel (NSW)


offers two major improvements to ATLAS muon spectrometry

rate capacity

detector elements much smaller in NSW than current SW

e.g. MM strip pitch  $\approx 0.5$  mm  
whereas CSC strip pitch  $\approx 5$  mm

smaller flux per element @ NSW



# the new Small Wheel (NSW)


offers two major improvements to ATLAS muon spectrometry

## trigger

NSW adds layer of coincidence for hardware trigger decision

powerful reduction of triggers not originating from muons

emphasis of this talk





# the new Small Wheel (NSW)


offers two major improvements to ATLAS muon spectrometry

trigger

NSW adds layer of coincidence for hardware trigger decision

powerful reduction of triggers not originating from muons


emphasis of this talk



# sTGC & MM triggers



# sTGC trigger electronics



American institutes building many pieces!


***BNL*** for FE chip: VMM

***Michigan*** for drivers: TDS and Router

***Illinois*** for trigger functions: Trigger anc.

# sTGC trigger electronics



- ask for many-layer coincidence with coarse pad triggers
- readout strips within pad trigger tower



large data reduction with pad “pre-triggering”




# sTGC trigger electronics



- 8 centroids of strips
- 2 centroids of centroids
- LUT for “track”

# MM trigger electronics



American institutes building many pieces!

**BNL** for FE chip: VMM

**Arizona** for FE card: MMFE8



**BNL** for driver card: ADDC

**Harvard** for trigger card: MM TP

**Illinois** for trigger functions: Trigger anc.

# MM trigger electronics


- only readout *first* strip per VMM  $\longrightarrow$  ~20x less data than full readout
- “ART”: Address in Real Time
- 64 strips per VMM covers ~2.5 cm



$\phi$  information inferred from stereo planes with  $\theta_{\text{strip}} = 1.5^\circ$



# MM trigger electronics




all ART strips received at  
MM TP, then filtered by  
roads in FPGA

large data reduction with  
road filtering:  
~500 roads per chamber

filtered strips then “fit”  
for track parameters

# MM trigger electronics




all ART strips received at  
MM TP, then filtered by  
roads in FPGA

large data reduction with  
road filtering:  
~500 roads per chamber

filtered strips then “fit”  
for track parameters

# MM trigger electronics



all ART strips received at  
MM TP, then filtered by  
roads in FPGA

large data reduction with  
road filtering:  
~500 roads per chamber

filtered strips then “fit”  
for track parameters


# performance

# simulated performance

simple OR of sTGC and MM decisions

excellent matching  
resolution between  
NSW and existing BW

NSW-BW coincidence  
efficiency expected to be  
90-95% or better for  
flagship L1  $p_T(\mu) > 20$  GeV






# simulated performance

simple OR of sTGC and MM decisions

muon trigger rates  
dominated by fakes  
in the endcaps


substantial rate  
reduction ( $\sim 3\times$ )  
predicted with NSW



# MM performance in data!



built 8-layer micromegas  
mini-chamber at Harvard to  
measure performance with  
cosmic ray muons

chamber equipped with  
prototype electronics,  
and recording both trigger  
and offline data



# MM cosmic ray test stand

taking offline (full) readout as reference,  
trigger algorithm fully functioning and performs within spec




# summary

# summary

New Small Wheel (NSW)  
is under construction now,  
planned to be operational for  
Run 3 of the LHC and beyond

NSW trigger hardware and  
algorithms in advanced state,  
already making triggers with  
cosmic ray muons





# bonus

# some references

New Small Wheel  
Technical Design Report

<https://atlas.web.cern.ch/Atlas/GROUPS/PHYSICS/UPGRADE/CERN-LHCC-2013-006/index.html>

Muon Spectrometer  
Technical Design Report

<http://atlas.web.cern.ch/Atlas/GROUPS/MUON/TDR/Web/TDR.html>

NSW Public Results

<https://twiki.cern.ch/twiki/bin/view/AtlasPublic/NSWPublicResults>

L1 Muon Public Results

<https://twiki.cern.ch/twiki/bin/view/AtlasPublic/L1MuonTriggerPublicResults>


Muon Trigger Public Results

<https://twiki.cern.ch/twiki/bin/view/AtlasPublic/MuonTriggerPublicResults>


# the current Small Wheel (SW)

innermost layer of ATLAS muon spectrometer endcap


not built to withstand HL-LHC!



## Monitored Drift Tubes (MDT)



## Cathode Strip Chambers (CSC)




# reminder: Harvard test stand

MMFE8s w/VMM2

ADDC V1

MMTP on VC707



collected a few 100k cosmic muons  
***with low noise*** in May

[Overview at last Muon Week](#)